CLASSIFICATION

SECRET

SECRET

REPORT

50X1-HUM

CENTRAL INTELLIGENCE AGENCY

INFORMATION FROM

FOREIGN DOCUMENTS OR RADIO BROADCASTS

CD NO.

DATE OF INFORMATION

1949

COUNTRY SUBJECT

: 57.54

Scientific - Chemistry, phosphorous

acid esters

DATE DIST. 6 Sep 1950

HOW **PUBLISHED** 

Bimonthly periodical

WHERE

USSR

**PUBLISHED** 

Mosc ow

NO. OF PAGES 3

DATE

PUBLISHED

May/Jun 1950

SUPPLEMENT TO

LANGUAGE

Russian

REPORT NO.

LUMENT CONTAINS INFORMATION AFFECTING THE MATIONAL DEFENSE UNITED STATES WITHIN THE MEANING OF ESPICIASE ACT S-21 AMD 28.4 AGENCES. ITS TRANSMISSION OF THE REVELATION CONTENTS IN ANY MANHEET OF AN UNAUTHORIZED FRANCH IS FRO ST LAW. REPRODUTION OF THIS FORM IS PROVIDED IN

THIS IS UNEVALUATED INFORMATION

SOURCE

, delta

Izvestiya Akademii Nauk SSSR, Otdeleniye Khimicheskikh Nauk, No 3, 1950.

## PARACHORS OF CYCLIC ESTERS OF CARBONIC, SULFUROUS, AND PHOSPHOROUS ACIDS

B. A. Arbuzov and V. S. Vinogradova Sci-Res Chem Inst imeni A. M. Butlerov, Kazan State U imeni V. I. Ul'yanov-Lenin Submitted 28 March 1949

The physical constants of the phosphorous acid esters listed below are of interest, because compounds of this general type may exhibit specific toxic properties or serve as intermediates in the preparation of toxic compounds. This refers to a potentiality rather than a. actual intention on the part of the Russian investigators. No actual intention to conduct work with the view of synthesizing toxic compounds has been mentioned in the article.

In order to determine the possibility of applying the group-correction method used in the case of five- and six-membered rings of carbocyclic compounds to the calculation of the parachors of heterocyclic compounds as well, the parachors of a number of cyclic esters of carbonic, sulfurous, and phosphorous acids were computed.

The group method (in which the parachors of individual groups of a ring are simply added) was found to be completely applicable for five- and six-membered heterocyclic rings containing an atom of oxygen, oxygen and sulfur, or oxygen and phosphorus in the ring.

The group values were determined for the parachors of the following groups contained in the rings:

		CLASSIFICATION	SECRET	
STATE	X NAVY	X NSRB	DISTRIBUTION	
ARMY	V AIR	X FB1		<u> </u>

Sanitized Copy Approved for Release 2011/08/17: CIA-RDP80-00809A000600340275-9

SECRET

SECRET

50X1-HUM

Cyclic Esters of Phosphorous Aciú

The author and his co-worker determined the parachors of four esters which V. M. Zoroastrova had previously (A. Ye. Arbuzov, V. M. Zoroastrova, and N. I. Rizpolozhenskiy, IAN SSSR, OKN, 208, 1948 / see, 00-W-4400/49/) synthesized through the action of alcohols on acid in the presence of pyr-synthesized through the action of alcohols on acid in the presence of pyr-idine or dimethylaniline. Her data for their constants are reproduced in the following table along with Arbuzov and Vinogradova's data on their parachors determined in this instance.

See table on following page

- 2 -

SECRET

SECRET

Sanitized Copy Approved for Release 2011/08/17 : CIA-RDP80-00809A000600340275-9

		Substance	<u>Bp</u>	d <sub>0</sub> 20	n <sub>D</sub> 20	<u>~<sup>20</sup></u>	Standard Ideal Value	Correction for Expansion	Parachor (computed)	Parachor (found)	Deviation in %	Corrections
		CH <sub>2</sub> - OP - OCH <sub>3</sub>	55-56°/ 23 mm	1.2200	1.4460	35.40	243.5	0.6	244.1	244.06	0	-5.4 for a five-membered ring
SEGRET		CH <sub>2</sub> - 0 P - OC <sub>2</sub> H <sub>5</sub>	51.5-52 <sup>0</sup> / 15 mm	1.1357	1.4398	38.47	281.5	0.7	282.2	281.4	-0.3	-5.4 -1.4 beta- correction for an ester chain
	secret	.5	62 <sup>0</sup> /13 mm	1.1102	1.4420	31.22	317.3	0.9	318.2	319.6	+0.5	-10.7 for a six-membered ring -3.2 for a side chain -1.7 me'ze-correction
		CH <sub>2</sub> - O CH <sub>2</sub> P - OC <sub>2</sub> H <sub>5</sub> CH - O	63-64°/ 8 mm	1.0709	1.4405	29.60	355•3	1.2	356.5	353•4	-0.8	-10.7 - 3.2 - 1.7 - 1.4

- F N D -

50X1-HUM

Sanitized Copy Approved for Release 2011/08/17 : CIA-RDP80-00809A000600340275-